

## Nucleophilic (Radio)Fluorination of Redox-Active Esters via Radical-Polar Crossover Enabled by Photoredox Catalysis

Eric W. Webb, John B. Park, Erin L. Cole, David J. Donnelly, Samuel J. Bonacorsi, William R. Ewing, and Abigail G. Doyle\*

Cite This: *J. Am. Chem. Soc.* 2020, 142, 9493–9500

Read Online

ACCESS |



Metrics &amp; More




Article Recommendations



Supporting Information

**ABSTRACT:** We report a redox-neutral method for nucleophilic fluorination of *N*-hydroxyphthalimide esters using an Ir photocatalyst under visible light irradiation. The method provides access to a broad range of aliphatic fluorides, including primary, secondary, and tertiary benzylic fluorides as well as unactivated tertiary fluorides, that are typically inaccessible by nucleophilic fluorination due to competing elimination. In addition, we show that the decarboxylative fluorination conditions are readily adapted to radiofluorination with [<sup>18</sup>F]KF. We propose that the reactions proceed by two electron transfers between the Ir catalyst and redox-active ester substrate to afford a carbocation intermediate that undergoes subsequent trapping by fluoride. Examples of trapping with O- and C-centered nucleophiles and deoxyfluorination via *N*-hydroxyphthalimidoyl oxalates are also presented, suggesting that this approach may offer a general blueprint for affecting redox-neutral S<sub>N</sub>1 substitutions under mild conditions.



## INTRODUCTION

Aliphatic organofluorine compounds are important structural motifs in pharmaceuticals, agrochemicals, and materials, conferring valuable biological and physical properties.<sup>1</sup> This motif is also prominently featured in positron emission tomography (PET) radiotracers.<sup>1a</sup> Consequently, the identification of mild methods for late-stage introduction of fluorine has been a longstanding goal, with distinct strategies arising using nucleophilic and electrophilic fluorine sources.<sup>2</sup> For reasons of cost, functional group compatibility, and translation to radiofluorination, researchers have sought mild methods for the preparation of fluoroalkyl groups using nucleophilic fluoride.<sup>1,2</sup> In most of these methods, Csp<sup>3</sup>–F bond formation follows a bimolecular nucleophilic substitution pathway (S<sub>N</sub>2 mechanism) and is thus limited to the preparation of activated or unhindered aliphatic fluorides (Scheme 1A).<sup>3</sup> Indeed, the typical restrictions on substrate scope for bimolecular nucleophilic substitution reactions are even more acute for fluoride due to its low nucleophilicity and high Brønsted basicity, leading to competitive elimination.<sup>4,5</sup>

Stepwise nucleophilic fluorination reactions that proceed through a carbocation intermediate (S<sub>N</sub>1 mechanism) provide a complementary strategy to bimolecular alkyl fluoride synthesis, enabling access to unactivated and hindered aliphatic fluorides. However, the generation of carbocation intermediates typically requires harsh Brønsted<sup>6</sup> or Lewis acidic conditions<sup>7</sup> that show poor functional group tolerance and lead to elimination and rearrangement pathways (Scheme 1B).

To overcome these limitations, researchers have recently explored new strategies for carbocation generation under nonacidic conditions. The Knowles group introduced a methodology to access carbocation intermediates via mesolytic cleavage following the oxidation of TEMPO-derived alkoxyamine substrates;<sup>8</sup> although a wide range of nucleophiles are compatible with this approach, the method was not shown to work with fluoride and the substrates can be challenging to access (Scheme 1C). Just recently, the Baran lab reported an electrochemical approach to carbocation generation from readily available carboxylic acids (Scheme 1D).<sup>9</sup> The method affords access to a broad range of hindered ethers from alcohol nucleophiles and four examples of alkylfluorides from KF. Although this report has significantly advanced the state of the art, the requirement for oxidizing conditions places limits on the substrate scope and the method was not shown to be amenable to radiofluorination. This report was closely followed by the disclosure of a photocatalytic decarboxylative ether synthesis by Ohmiya, Nagao, and co-workers.<sup>10</sup> Yet further discovery and development of complementary methods for carbocation generation are necessary to enable broad access to

Received: March 20, 2020

Published: May 7, 2020




ACS Publications

© 2020 American Chemical Society


9493

<https://dx.doi.org/10.1021/jacs.0c03125>  
*J. Am. Chem. Soc.* 2020, 142, 9493–9500


## Scheme 1. Nucleophilic Fluorination

A. Bimolecular nucleophilic substitution ( $S_N2$ ) with fluoride

- Limited to 1° & 2° aliphatic fluorides; 3° inaccessible
- Elimination competitive

B. Lewis or Brønsted acid-facilitated  $S_N1$ 


- Caustic reagents preclude late-stage functionalization
- Access to 3° fluorides

C. Photocatalyzed  $S_N1$  via mesolytic cleavage (Knowles)

- C-C, C-N, C-O bond formation

D. Electrochemical oxidation  $S_N1$  (Baran)

- 4 examples

E. This work: Redox-neutral, radical-polar crossover  $S_N1$  substitution

- Redox neutral
- Broad FG tolerance and mild conditions
- Access to 1°, 2°, 3° fluorides

hindered and unactivated aliphatic fluorides, including for late-stage fluorination and radiofluorination, as well as expanding the repertoire for additional synthetic applications.

In this context, we sought to develop a redox-neutral method for carbocation generation via radical-polar crossover<sup>10,11</sup> from *N*-hydroxyphthalimide esters (Scheme 1 E). We selected redox active esters as a substrate class since numerous research groups have recently reported their use as precursors to alkyl radicals via single-electron reduction and decarboxylation with release of a non-nucleophilic leaving group.<sup>12</sup> We hypothesized that upon single-electron reduction of the substrate by the excited state of a suitable photoredox catalyst, the resulting radical could be oxidized to the corresponding carbocation by the oxidized photocatalyst and subsequently trapped with fluoride.<sup>13</sup> Given the similar trends in stability between radical and carbocation species,<sup>14</sup> we anticipated that highly substituted aliphatic substrates should be particularly amenable to both radical and carbocation generation in a radical-polar crossover, thereby expanding the scope of aliphatic fluorides available via nucleophilic fluorination. This approach would also provide a useful complement to radical methodologies for decarboxylative fluorination that require oxidizing electrophilic fluorine sources<sup>15</sup> or the combination of a nucleophilic fluoride source with a stoichiometric oxidant, as described by the Groves lab.<sup>16</sup>

Here we report a redox-neutral decarboxylative nucleophilic fluorination that delivers primary, secondary, and tertiary benzylic fluorides and unactivated tertiary fluorides with broad functional group tolerance. We also describe mechanistic experiments that provide evidence for both radical and carbocation intermediates; in so doing, we present applications


to the construction of sterically congested ethers and C-C bonds, establishing the generality of the strategy as a blueprint for affecting redox-neutral  $S_N1$  substitutions. Finally, we show that the method is amenable to <sup>18</sup>F-radiofluorination.

## RESULTS AND DISCUSSION

**Optimization.** We initiated our studies with the *N*-hydroxyphthalimide ester **1** derived from naproxen, as naproxen's electron-rich arene would likely be incompatible with decarboxylative fluorination conditions that utilize electrophilic fluorine sources or stoichiometric oxidants.<sup>15b</sup> Subjecting **1** to irradiation with 34 W blue LEDs in the presence of 1 mol % Ir(dF-ppy)<sub>3</sub> **3** and 3 equiv of Et<sub>3</sub>N·3HF delivered benzylic fluoride **2** in almost quantitative yield (Table 1, entry 1). Fluorination does not proceed in the

Table 1. Optimization of Reaction Conditions<sup>a,b</sup>


entry	deviation from standard conditions	yield 2 (%) <sup>c</sup>
1	No deviation	99
2	No photocatalyst	0
3	No light	0
4	Ir(ppy) <sub>3</sub> <b>4</b> instead of <b>3</b>	96
5	[Ir(dF-CF <sub>3</sub> -ppy) <sub>2</sub> (dtbpy)]PF <sub>6</sub> <b>5</b> instead of <b>3</b>	94
6	4CzIPN <b>6</b> instead of <b>3</b>	91
7	THF instead of DCM	22
8	MeCN instead of DCM	68
9	3 equiv. KF instead of Et <sub>3</sub> N·3HF	0
10	3 equiv. KF/10 equiv. HFIP instead of Et <sub>3</sub> N·3HF	4
11	4 mmol scale <sup>d</sup>	90 (75) <sup>e</sup>
12	0.0625 mol% Ir(dF-ppy) <sub>3</sub>	85




<sup>a</sup>0.2 mmol scale. <sup>b</sup>All potentials given are versus a saturated calomel electrode (SCE) and taken from ref 19. <sup>c</sup>Yields determined by <sup>19</sup>F-NMR using 1-fluoronaphthalene as an external standard. <sup>d</sup>General conditions except 0.4 M, 1.5 equiv of Et<sub>3</sub>N·3HF. <sup>e</sup>Isolated yield.


absence of light or photocatalyst, resulting in recovery of starting material (Table 1, entries 2 and 3). More reducing or oxidizing iridium photocatalysts were competent in the reaction, as was the organic photocatalyst 4CzIPN, all displaying high yet diminished reactivity compared to photocatalyst **3** (Table 1, entries 4–6). While DCM was found to be the optimal solvent for the reaction, fluorination proceeded with moderate to good yield in tetrahydrofuran and

Table 2. Substrate Scope for Photocatalytic Decarboxylative Fluorination<sup>a</sup>








  
 phthalimide ester

**3** (1 mol%)  
 Et<sub>3</sub>N·3HF (3 equiv)  
 34 W Blue LED  
 DCM (0.2 M), rt, 4–6 h






  
 aliphatic fluoride  
 7–40

  
**3**






### Primary and secondary benzylic

 <b>7–14</b>	<b>7</b> - <i>t</i> -Bu <b>64% yield</b> <b>8</b> -Ph <b>59% yield</b> <b>9</b> -I <b>60% yield</b> <b>10</b> -OBn <b>76% yield</b>	<b>11</b> -SMe <b>62% yield<sup>b</sup></b> <b>12</b> -NHAc <b>73% yield</b> <b>13</b> -OCF <sub>3</sub> <b>60% yield<sup>b</sup></b> <b>14</b> -CF <sub>3</sub> <b>6% yield<sup>b</sup></b>
 <b>15</b> <b>63% yield</b>	 <b>16</b> <b>40% yield<sup>b</sup></b>	 <b>17</b> <b>93% yield</b>
 <b>19</b> <b>64% yield</b>	 <b>20</b> <b>56% yield</b>	 <b>21</b> <b>65% yield</b> <i>from ketoprofen</i>




### α-Heteroatom

 <b>32</b> <b>28% yield<sup>b</sup></b>	 <b>33</b> <b>53% yield</b>	 <b>34</b> <b>44% yield<sup>d</sup></b>
 <b>35</b> <b>57% yield<sup>b,e</sup></b> <i>from trillipix</i>	 <b>36</b> <b>56% yield</b> <i>from cyhalofop</i>	




### Tertiary benzylic and aliphatic

 <b>22</b> <b>87% yield</b>	<p> <b>23</b> <b>3</b> <b>92% yield</b>  <b>24</b> <b>2</b> <b>85% yield<sup>b,e</sup></b>  <b>25</b> <b>1</b> <b>90% yield</b> </p>	 <b>26</b> <b>70% yield<sup>b</sup></b>	 <b>27</b> <b>40% yield<sup>b</sup></b>
<p> <b>28</b> -<i>t</i>-Bu <b>76% yield<sup>b</sup></b>  <b>29</b> -CN <b>50% yield</b> </p>	 <b>30</b> <b>66% yield</b> <i>from gemfibrozil</i>	 <b>31</b> <b>32% yield<sup>b,c</sup></b> <i>from enoxalone</i>	

### Multifluorinated

 <b>37</b> <b>56% yield</b>	 <b>38</b> <b>89% yield<sup>b</sup></b> <i>from Mosher's acid</i>	 <b>39</b> <b>52% yield</b>
--	--	---

### Tertiary deoxyfluorination

 <b>22</b> <b>55% yield<sup>b,f</sup></b>	 <b>40</b> <b>33% yield<sup>b</sup></b>	<p>from:</p> 
--	--	--

<sup>a</sup>Isolated yields of an average of two runs on 0.5–1.0 mmol scale. <sup>b</sup>Yield determined by <sup>19</sup>F-NMR using 1-fluoronaphthalene as an external standard. <sup>c</sup>1.3:1 d.r. <sup>d</sup>>20:1 d.r. <sup>e</sup>Product unstable. <sup>f</sup>Yield corrected for starting material contamination with residual alcohol.

acetonitrile, solvents that are commonly used in radiofluorination (Table 1, entries 7 and 8). Only trace product was observed using KF/HFIP (1,1,1,3,3,3-hexafluoroisopropanol)<sup>17</sup> instead of Et<sub>3</sub>N·3HF, and no product was observed using KF in the absence of HFIP (Table 1, entries 9 and 10). Although impractical from a preparative standpoint, this result provides support for possible translation to radiofluorination since [<sup>18</sup>F]KF is the most common reagent for <sup>18</sup>F-radiochemistry (*vide infra*).<sup>18</sup> Finally, we found that the reaction could be readily scaled to 4 mmol (Table 1, entry 11) and photocatalyst loading could be reduced to 0.0625 mol % with minimal impact on the reaction efficiency (Table 1, entry 12). This speaks to the practicality of the method, and the ability to use such low photocatalyst loading holds important mechanistic implications (*vide infra*).

**Scope Elucidation.** With optimized conditions in hand, we evaluated the scope of the transformation. We found that a variety of primary benzylic fluorides could be obtained in good to excellent yields (Table 2, 7–13, 15, and 16). Several substrates bearing electron-rich functionality otherwise suscep-

tible to oxidation under previously reported decarboxylative fluorination conditions were tolerated,<sup>15b</sup> delivering dioxole 15, benzyl-protected phenol 10, and thioether 11. Fluorinated products bearing medically relevant amides and trifluoromethoxy groups were generated in good yields (12 and 13). Whereas these electron-rich and electron-neutral primary benzylic phthalimide esters were competent substrates, electron-deficient substrates afforded benzylic fluorides in low yield at high conversion (14), presumably because single-electron reduction and decarboxylation to the carbon-centered radical is facile but oxidation of the radical to the cation is disfavored due to the electron-withdrawing substituent.<sup>13</sup> As expected on the basis of this hypothesis, we found that replacing the primary benzylic substrate with a secondary substrate bearing the same substitution pattern restored reactivity, with fluorinated product 17 obtained in 93% yield.

A common limitation in nucleophilic fluorination methods that deliver secondary benzylic fluorides is elimination to styrene byproducts.<sup>3d,f</sup> For all of the secondary substrates

examined in Table 2, less than 5% elimination was observed, a testament to the mildness of the conditions. Indeed, even product **19** was obtained in high yield with minimal elimination despite the presence of a  $\beta$ -carbonyl functional group. By comparison, access to  $\beta$ -fluoro carbonyl derivatives by deoxyfluorination has presented a major challenge to date due to competing elimination.<sup>3d,f</sup> Several handles for subsequent transition metal-mediated coupling were tolerated, including aryl iodides **9**, bromides **16** and **33**, and chlorides **18** and **23–26**. This tolerance for easily reduced functionality can even be extended to azide-containing product **20**, which was generated in 56% yield and offers a reactive handle for subsequent “click” chemistry that is widely used in bioconjugation.<sup>21</sup> Additionally, basic heterocycles and hetero-aromatic groups otherwise susceptible to oxidation or Minisci chemistry underwent decarboxylative fluorination in good yields (**18** and **26**).

In contrast to typical methods for nucleophilic fluorination, we found that access to benzylic and unactivated tertiary fluorides is possible. For example, acyclic **22** as well as tertiary benzylic fluorides embedded within carbocyclic and heterocyclic ring systems are generated in 70–92% yield, as in the cases of **23–26**. Likewise, both cyclic and acyclic unactivated tertiary fluorides could be obtained (**27–31**). Whereas neighboring group participation may be operative in the generation of the homobenzylic tertiary fluorides **28** and **29**, it does not appear to be necessary given the success of the cyano-substituted homobenzylic fluoride **29** and fluorides **27** and **31** that do not possess a proximal nucleophilic residue. Notably, we were able to extend this protocol to the fluorination of gemfibrozil **30** in 66% yield.

Fluoroether and fluorothioether functionality has been shown to confer unique and valuable properties to biologically active small molecules.<sup>22</sup> We found that the redox-neutral decarboxylative nucleophilic fluorination also delivers  $\alpha$ -oxy- and  $\alpha$ -thioether motifs in moderate to good yield (**32** and **33**). As a demonstration of the viability of the method for late-stage derivatization, fluorinated ribose **34**, trillipix-derivative **35**, and the herbicide cyhalofop-derived **36** were all readily accommodated. Likewise, application of the optimal conditions to the preparation of difluoromethyl and perfluorinated groups was successful,<sup>23</sup> as in the cases of **37** and **38**, and permitted the synthesis of difluorofluorene **39**, a motif featured in the hepatitis C drug ledipasvir.

Fluorine incorporation is commonly used as a bioisostere for several functionalities including C–OH and C–H bonds.<sup>1b</sup> In this regard, the conversion of abundant alcohols into the corresponding alkyl fluorides via deoxyfluorination represents an attractive synthetic disconnection. However, deoxyfluorinations of tertiary alcohols to access tertiary fluorides are typically unsuccessful.<sup>24</sup> MacMillan and co-workers have recently reported a deoxyfluorination of oxalate half-esters to access tertiary fluorides.<sup>25</sup> However, the method uses an electrophilic fluorine source. Since *tert*-alkyl *N*-hydroxyphthalimidoyl oxalates are similar in redox potential to *N*-hydroxyphthalimide esters, we hypothesized that these may be amenable to the catalytic nucleophilic fluorination strategy outlined herein.<sup>26</sup> Indeed, we were pleased to find that tertiary fluorides **22** and **40** could be obtained in 55% and 33% yield from *tert*-alkyl *N*-hydroxyphthalimidoyl oxalate esters under otherwise identical conditions. Since these substrates are readily available from alcohols, the method represents a complementary approach to

nucleophilic deoxyfluorination which is typically limited to primary and secondary alcohols.<sup>3a–f</sup>

**Mechanistic Investigations.** We propose that excited **3** ( $E_{1/2}^* = -1.28$  vs SCE<sup>19a</sup>) undergoes single-electron transfer (SET) with the *N*-hydroxyphthalimide ester **A** ( $\sim -1.3$  V vs SCE<sup>20</sup>) (Figure 1). Fragmentation of the resulting phthalimide





Figure 1. Mechanistic proposal.

ester radical anion and subsequent extrusion of carbon dioxide generate carbon-centered radical **B**. Radical intermediate **B** ( $E_{1/2}^{\text{ox}} = < 0.73$  V vs SCE for 1° benzylic,  $E_{1/2}^{\text{ox}} = 0.09$  V vs SCE for tertiary aliphatic<sup>13b</sup>) is then oxidized by photocatalyst **3** ( $\text{Ir}^{\text{IV}}/\text{Ir}^{\text{III}} E_{1/2} = 0.94$  V vs SCE<sup>19a</sup>), turning over the photocatalyst and furnishing carbocation **C**. Finally, this carbocation is trapped by the fluoride source to furnish the desired alkyl fluoride **D**.

A number of experimental observations provide support for the proposed mechanism. Stern–Volmer quenching analysis of the individual components of the reaction mixture indicates that the phthalimide ester quenches the excited state photocatalyst with an observed  $K_{\text{SV}}$  of  $4.7 \times 10^9 \text{ M}^{-1} \text{ s}^{-1}$ , which is similar to the quenching rate of the reaction mixture.<sup>27</sup> The quantum yield of this fluorination reaction is 0.37, indicating that chain mechanisms are unlikely or inefficient.<sup>27,28</sup> This result, combined with the observation that the fluorinations proceed with high reaction efficiency at extremely low photocatalyst loadings (0.0625 mol % of **3** in Table 1, entry 12), suggests that the reaction is unimolecular in photocatalyst.

Subjecting tertiary phthalimide ester **41** to the fluorination conditions in the presence of several known radical traps provides evidence for the intermediacy of a radical. For example, addition of TEMPO (2,2,6,6-tetramethyl-1-piperidin-1-yl) to the reaction resulted in complete inhibition of fluorination, with concomitant detection of TEMPO adduct **42** (Figure 2). In the presence of methyl acrylate, radical addition product **43** was observed, along with a diminished yield of tertiary fluoride **30** (66% isolated vs 45% by  $^{19}\text{F}$  NMR in the presence of methyl acrylate). However, fluoride is not incorporated into **43**, presumably because radical oxidation adjacent to the ester carbonyl is unfavorable. On the other hand, conducting the fluorination reaction in the presence of 1.5 equiv of styrene afforded a new fluorinated product **44** in addition to the direct fluorination product **30**. **44** is most likely





**Figure 2.** Radical and cation trapping. <sup>a</sup>General conditions: 1 mol % Ir(dF-ppy)<sub>3</sub>, 0.2 mmol of **41**, 3 equiv of Et<sub>3</sub>N·3HF. <sup>b</sup>1.5 equiv of TEMPO. <sup>c</sup>1.5 equiv of methyl acrylate. <sup>d</sup>1.5 equiv of styrene. <sup>e</sup>3 equiv of 1,3,5-trimethoxybenzene and 0.3 equiv of Et<sub>3</sub>N·3HF. <sup>f</sup>5 equiv of phenol and 0.3 equiv of Et<sub>3</sub>N·3HF. <sup>g</sup>6 equiv of methanol and 0.3 equiv of Et<sub>3</sub>N·3HF. <sup>h</sup>5 equiv of HFIP and run for 24 h; <sup>19</sup>F NMR yield vs external 1-fluoronaphthalene.

generated via addition of radical **B** to the styrene followed by oxidation of the resulting benzylic radical and trapping with fluoride. Evaluation of a series of electronically differentiated styrene substrates, and the outcome of the reaction with methyl acrylate, provides evidence against an alternative pathway wherein radical oxidation to cation **C** precedes olefin addition.<sup>27</sup>

As a test for the intermediacy of a carbocation, we investigated whether other polar nucleophiles could be used in the decarboxylative substitution reaction. Notably, we found that C- and O-centered nucleophiles were competent with only minor changes to the reaction parameters (Table 3, 45–48).<sup>29</sup> For example, 1,3,5-trimethoxybenzene underwent addition to generate benzhydryl **45** in 81% yield. This reaction represents a Friedel–Crafts substitution without a Lewis acid using abundant carboxylic acid precursors in place of alkyl halide substrates. Likewise, several alcohol nucleophiles delivered ether products **46–48** in good yield. The broad tolerance for a range of nucleophiles, including sterically hindered and poorly nucleophilic species, implicates the intermediacy of a carbocation and demonstrates the generality of the strategy to effect a range of challenging substitution reactions under remarkably mild conditions.

**Radiochemistry.** Aliphatic <sup>18</sup>F-radiolabeled PET tracers are almost exclusively prepared by nucleophilic substitution of

**Table 3.** Development of Radiochemical Protocol<sup>a</sup>

Entry	Deviation from Standard Conditions	RCC (%) <sup>b</sup>
1	none	64 <sub>n=2</sub>
2	no photocatalyst	0 <sub>n=1</sub>
3	no light	0 <sub>n=2</sub>
4	<b>3</b> instead of Ir(F-ppy) <sub>3</sub>	6 <sub>n=2</sub>
5	Ir(CF <sub>3</sub> -ppy) <sub>3</sub> instead of Ir(F-ppy) <sub>3</sub>	43 <sub>n=2</sub>
6	Ir(dF- <i>t</i> -Bu-ppy) <sub>3</sub> instead of Ir(F-ppy) <sub>3</sub>	24 <sub>n=2</sub>
7	Ir(F- <i>t</i> -Bu-ppy) <sub>3</sub> instead of Ir(F-ppy) <sub>3</sub>	49 <sub>n=2</sub>
8	2 min instead of 10 min	62 <sub>n=1</sub>
9	100 μL HFIP added	33 <sub>n=2</sub>

<sup>a</sup>Typical reaction conditions: 5.3 mol of **1**, 16 mol % photocatalyst, 0.7 mL of MeCN, and [<sup>18</sup>F]F<sup>−</sup>/K<sub>2</sub>CO<sub>3</sub>/K<sub>222</sub> (~0.370 GBq of activity per reaction). <sup>b</sup>RCC was determined by radio-TLC with number of replicates noted.


alkyl sulfonates with [<sup>18</sup>F]KF in the presence of a phase transfer reagent Kryptofix 2.2.2 (K<sub>222</sub>).<sup>18</sup> As such, access to high specific activity radiolabeled targets bearing unactivated secondary or tertiary fluorides remains a critical challenge. Moreover, the harsh conditions necessary for substitution of even primary or activated secondary substrates (>100 °C and high basicity) are often not suitable for late-stage radiofluorination and often lead to inseparable olefin byproducts.<sup>18</sup> We anticipated that successful translation of the photocatalytic decarboxylative nucleophilic fluorination method would therefore enable access to previously challenging or impossible to prepare radiotracers.

To translate our method, we elected to pursue the decarboxylative radiofluorination in acetonitrile to avoid any potential clinical issues with dichloromethane. Testing the radiofluorinations in CH<sub>3</sub>CN using [<sup>18</sup>F]KF/K<sub>222</sub> as the fluoride source, we found that the previously optimized photocatalyst **3** was no longer the most effective, with Ir(F-ppy)<sub>3</sub> instead affording the highest radiochemical incorporation (Table 3, entries 1 and 4–7).<sup>27</sup> No radiofluorination was observed in the absence of light or photocatalyst as determined by radio-TLC or radio-HPLC (Table 3, entries 2 and 3). While we had previously found that HFIP was necessary for achieving fluorination using [<sup>19</sup>F]KF under the nonradiochemical conditions (Table 1, entries 10 and 11), the addition of HFIP proved detrimental in the radiochemical system (Table 3, entry 9). Under the optimal conditions, the reaction is particularly fast, furnishing [<sup>18</sup>F]**2** in 62% radiochemical conversion (RCC) within 2 min of irradiation (Table 3, entry 8). The radiofluorination was also scalable (1850–3700 MBq) albeit with diminished radiochemical conversion, permitting generation of sufficient quantities of [<sup>18</sup>F]**2** to determine its molar activity (36.6 ± 18.8 GBq/μmol, decay corrected to end of synthesis), which is on par with other no-carrier added nucleophilic fluorination protocols.<sup>17,18,30</sup> Notably, these radiochemical reactions were conducted in a 3D-printed apparatus that permits automated

handling of radioactivity with irradiation taking place from the bottom of the vials.<sup>27</sup>

Access to [<sup>18</sup>F]**2** is significant since its synthesis by S<sub>N</sub>2 displacement with [<sup>18</sup>F]KF would likely be plagued by rapid formation of elimination byproduct. We also briefly explored application of the radiofluorination conditions to the radio-synthesis of other fluorinated motifs that are challenging to access using conventional methods (Table 4). For example, the

**Table 4. Scope of Photocatalyzed Radiofluorination**



<sup>a</sup>Typical reaction conditions: 5.3 mol of **1**, 0.6 mg of Ir(F-ppy)<sub>3</sub>, 0.7 mL of MeCN, and [<sup>18</sup>F]F<sup>−</sup>/K<sub>2</sub>CO<sub>3</sub>/K<sub>222</sub> (~0.370 GBq of activity per reaction). RCC was determined by radio-TLC with number of replicates noted. Product identity was typically confirmed by HPLC coinjection. <sup>b</sup>Reactions were conducted with 1.85–3.70 GBq of activity per reaction. <sup>c</sup>Reactions conducted using photocatalyst **3**. <sup>d</sup>Product identity confirmed via TLC.

radiofluorination protocol is amenable to the installation of a tertiary fluoride: [<sup>18</sup>F]**30** derived from gemfibrozil was obtained in 9 ± 2% RCC. Furthermore, we found that ribose analogue [<sup>18</sup>F]**34** could be prepared in 42% RCC. In this case, the sulfonate precursor readily decomposes at room temperature, severely limiting access to radiolabeled ribose analogues by conventional substitution reactions.<sup>31</sup>

## CONCLUSION

We have developed a photocatalytic method for nucleophilic fluorination of *N*-hydroxyphthalimide esters that exploits the redox activity of radicals as a route to carbocation formation. The approach generates a variety of useful fluorinated motifs under mild conditions and is compatible with functional groups that challenge other synthetic methods using both nucleophilic and electrophilic fluorine sources, such as access to tertiary aliphatic fluorides and tolerance to electron-rich functionality. Moreover, translation of the method to a radiochemical protocol was possible, enabling radiofluorination of derivatives of bioactive molecules. We present a preliminary demonstration of the generality of this approach to redox-neutral S<sub>N</sub>1-like substitutions in extensions to a new substrate class, such as deoxyfluorination of tertiary *N*-hydroxyphthalimidoyl oxalates, and to new nucleophiles, as in the construction of sterically congested ethers and C–C bonds.

## ASSOCIATED CONTENT

### Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/jacs.0c03125>.

Experimental procedures and characterization and spectral data for new compounds (PDF)

## AUTHOR INFORMATION

### Corresponding Author

Abigail G. Doyle – Department of Chemistry, Princeton University, Princeton, New Jersey 08544, United States; [orcid.org/0000-0002-6641-0833](https://orcid.org/0000-0002-6641-0833); Email: [agdoyle@princeton.edu](mailto:agdoyle@princeton.edu)

### Authors

Eric W. Webb – Department of Chemistry, Princeton University, Princeton, New Jersey 08544, United States  
John B. Park – Department of Chemistry, Princeton University, Princeton, New Jersey 08544, United States  
Erin L. Cole – Discovery Chemistry Platforms, PET Radiochemical Synthesis, Bristol-Myers Squibb Research and Development, Princeton, New Jersey 08543, United States  
David J. Donnelly – Discovery Chemistry Platforms, PET Radiochemical Synthesis, Bristol-Myers Squibb Research and Development, Princeton, New Jersey 08543, United States  
Samuel J. Bonacorsi – Discovery Chemistry Platforms, PET Radiochemical Synthesis, Bristol-Myers Squibb Research and Development, Princeton, New Jersey 08543, United States  
William R. Ewing – Discovery Chemistry, Bristol-Myers Squibb, Princeton, New Jersey 08543-5400, United States

Complete contact information is available at: <https://pubs.acs.org/doi/10.1021/jacs.0c03125>

### Notes

The authors declare no competing financial interest.

## ACKNOWLEDGMENTS

Financial support was generously provided by NSF (CHE-1565983) and Bristol-Myers Squibb. We thank Dhar Murali (BMS) for helpful suggestions, Erin E. Gomes (BMS) for radiochemical and analytical expertise, and Daniel Batella (BMS) for construction of the apparatus to allow radiochemical irradiation. István Pelzer, Kenneth Conover, and John Eng are acknowledged for analytical aid. We thank Laura K. G. Ackermann, Meredith A. Borden, and Talia J. Steiman for helpful discussions.

## REFERENCES

- (1) (a) Gillis, E. P.; Eastman, K. J.; Hill, M. D.; Donnelly, D. J.; Meanwell, N. A. Applications of Fluorine in Medicinal Chemistry. *J. Med. Chem.* **2015**, *58*, 8315. (b) Jeschke, P. The Unique Role of Fluorine in the Design of Active Ingredients for Modern Crop Protection. *ChemBioChem* **2004**, *5*, 570. (c) Hagmann, W. K. The Many Roles for Fluorine in Medicinal Chemistry. *J. Med. Chem.* **2008**, *51*, 4359.
- (2) (a) Liang, T.; Neumann, C. N.; Ritter, T. Introduction of Fluorine and Fluorine-Containing Functional Groups. *Angew. Chem., Int. Ed.* **2013**, *52*, 8214. (b) Neumann, C. N.; Ritter, T. Late-Stage Fluorination: Fancy Novelty or Useful Tool? *Angew. Chem., Int. Ed.* **2015**, *54*, 3216.
- (3) Deoxyfluorination with DAST: (a) Middleton, W. J. New Fluorinating Reagents. Dialkylaminosulfur Fluorides. *J. Org. Chem.* **1975**, *40*, 574. (b) Markovskij, L. N.; Pashinnik, V. E.; Kirsanov, A. V. Application of Dialkylaminosulfur Trifluorides in the Synthesis of Fluoroorganic Compounds. *Synthesis* **1973**, *1973*, 787. Deoxyfluorination with Phenofluor/Alkylfluor: (c) Tang, P.; Wang, W.; Ritter, T. Deoxyfluorination of Phenols. *J. Am. Chem. Soc.* **2011**, *133*, 11482. (d) Sladojevich, F.; Arlow, S. I.; Tang, P.; Ritter, T. Late-Stage

- Deoxyfluorination of Alcohols with Phenofluor. *J. Am. Chem. Soc.* **2013**, *135*, 2470. (e) Goldberg, N. W.; Shen, X.; Li, J.; Ritter, T. AlkylFluor: Deoxyfluorination of Alcohols. *Org. Lett.* **2016**, *18*, 6102. Deoxyfluorination with Pyfluor: (f) Nielsen, M. K.; Ugaz, C. R.; Li, W.; Doyle, A. G. PyFluor: A Low-Cost, Stable, and Selective Deoxyfluorination Reagent. *J. Am. Chem. Soc.* **2015**, *137*, 9571. Fluorination with hypervalent iodine reagents: (g) Hara, S.; Nakahigashi, J.; Ishi-i, K.; Sawaguchi, M.; Sakai, H.; Fukuhara, T.; Yoneda, N. Difluorination of Alkenes with Iodotoluene Difluoride. *Synlett* **1998**, 1998, 495. (h) Chen, H.; Kaga, A.; Chiba, S. Anti-Selective Aminofluorination of Alkenes with Amidines Mediated by Hypervalent Iodine(III) Reagents. *Org. Biomol. Chem.* **2016**, *14*, 5481. (i) Kong, W.; Feige, P.; De Haro, T.; Nevado, C. Regio- and Enantioselective Aminofluorination of Alkenes. *Angew. Chem., Int. Ed.* **2013**, *52*, 2469. (j) Mennie, K. M.; Banik, S. M.; Reichert, E. C.; Jacobsen, E. N. Catalytic Diastereo- and Enantioselective Difluorination of Alkenes. *J. Am. Chem. Soc.* **2018**, *140*, 4797. (k) Banik, S. M.; Medley, J. W.; Jacobsen, E. N. Catalytic, Asymmetric Difluorination of Alkenes to Generate Difluoromethylated Stereocenters. *Science* **2016**, *353*, 51. (l) Banik, S. M.; Medley, J. W.; Jacobsen, E. N. Catalytic, Diastereoselective 1,2-Di Fluorination of Alkenes. *J. Am. Chem. Soc.* **2016**, *138*, 5000. Fluorination via ion-binding catalysis: (m) Pupo, G.; Ibba, F.; Ascough, D. M. H. H.; Vicini, A. C.; Ricci, P.; Christensen, K. E.; Pfeifer, L.; Morphy, J. R.; Brown, J. M.; Paton, R. S.; Gouverneur, V. Asymmetric Nucleophilic Fluorination under Hydrogen Bonding Phase-Transfer Catalysis. *Science* **2018**, *360*, 638. (n) Pupo, G.; Vicini, A. C.; Ascough, D. M. H.; Ibba, F.; Christensen, K. E.; Thompson, A. L.; Brown, J. M.; Paton, R. S.; Gouverneur, V. Hydrogen Bonding Phase-Transfer Catalysis with Potassium Fluoride: Enantioselective Synthesis of  $\beta$ -Fluoroamines. *J. Am. Chem. Soc.* **2019**, *141*, 2878. (o) Pfeifer, L.; Engle, K. M.; Pidgeon, G. W.; Sparkes, H. A.; Thompson, A. L.; Brown, J. M.; Gouverneur, V. Hydrogen-Bonded Homoleptic Fluoride-Diaryurea Complexes: Structure, Reactivity, and Coordinating Power. *J. Am. Chem. Soc.* **2016**, *138*, 13314.
- (4) (a) Liang, S.; Hammond, G. B.; Xu, B. Hydrogen Bonding: Regulator for Nucleophilic Fluorination. *Chem. - Eur. J.* **2017**, *23*, 17850. (b) Engle, K. M.; Pfeifer, L.; Pidgeon, G. W.; Giuffredi, G. T.; Thompson, A. L.; Paton, R. S.; Brown, J. M.; Gouverneur, V. Coordination Diversity in Hydrogen-Bonded Homoleptic Fluoride-Alcohol Complexes Modulates Reactivity. *Chem. Sci.* **2015**, *6*, 5293. (c) Nolte, C.; Ammer, J.; Mayr, H. Nucleofugality and Nucleophilicity of Fluoride in Protic Solvents. *J. Org. Chem.* **2012**, *77*, 3325. (d) Landini, D.; Maia, A.; Rampoldi, A. Dramatic Effect of the Specific Solvation on the Reactivity of Quaternary Ammonium Fluorides and Poly(Hydrogen Fluorides),  $(\text{HF})_n \bullet \text{F}^-$ , in Media of Low Polarity. *J. Org. Chem.* **1989**, *54*, 328. (e) Lee, J.-W.; Oliveira, M. T.; Jang, H. B.; Lee, S.; Chi, D. Y.; Kim, D. W.; Song, C. E. Hydrogen-Bond Promoted Nucleophilic Fluorination: Concept, Mechanism and Applications in Positron Emission Tomography. *Chem. Soc. Rev.* **2016**, *45*, 4638.
- (5) Liotta, C. L.; Harris, H. P. The Chemistry of "Naked" Anions. I. Reactions of the 18-Crown-6 Complex of Potassium Fluoride with Organic Substrates in Aprotic Organic Solvents. *J. Am. Chem. Soc.* **1974**, *96*, 2250.
- (6) (a) Olah, G. A.; Nojima, M.; Kerekes, I. Hydrofluorination of Alkenes, Cyclopropane, and Alkynes with Poly-Hydrogen Fluoride/Pyridine (Trialkylamine) Reagents. *Synthesis* **1973**, 1973, 779. (b) Olah, G. A.; Welch, J. T.; Vankar, Y. D.; Nojima, M.; Kerekes, I.; Olah, J. A. Pyridinium Poly(Hydrogen Fluoride) (30% Pyridine-70% Hydrogen Fluoride): A Convenient Reagent for Organic Fluorination Reactions. *J. Org. Chem.* **1979**, *44*, 3872. (c) Lu, Z.; Zeng, X.; Hammond, G. B.; Xu, B. Widely Applicable Hydrofluorination of Alkenes via Bifunctional Activation of Hydrogen Fluoride. *J. Am. Chem. Soc.* **2017**, *139*, 18202. (d) Bertrand, X.; Paquin, J.-F. Direct Hydrofluorination of Methallyl Alkenes Using a Methanesulfonic Acid/Triethylamine Trihydrofluoride Combination. *Org. Lett.* **2019**, *21*, 9759.
- (7) Silver mediated fluorination: (a) Khotavivattana, T.; Verhoog, S.; Tredwell, M.; Pfeifer, L.; Calderwood, S.; Wheelhouse, K.; Leecollier, T.; Gouverneur, V.  $^{18}\text{F}$ -Labeling of Aryl- $\text{SCF}_3$ ,  $\text{-OCF}_3$  and  $\text{-OCHF}_2$  with  $^{18}\text{F}$  Fluoride. *Angew. Chem., Int. Ed.* **2015**, *54*, 9991. (b) Verhoog, S.; Pfeifer, L.; Khotavivattana, T.; Calderwood, S.; Collier, T. L.; Wheelhouse, K.; Tredwell, M.; Gouverneur, V. Silver-Mediated  $^{18}\text{F}$ -Labeling of Aryl- $\text{CF}_3$  and Aryl- $\text{CHF}_2$  with  $^{18}\text{F}$ -Fluoride. *Synlett* **2015**, 27, 25. Boron trifluoride mediated fluorination and references therein: (c) Cresswell, A. J.; Davies, S. G.; Roberts, P. M.; Thomson, J. E. Beyond the Balz-Schiemann Reaction: The Utility of Tetrafluoroborates and Boron Trifluoride as Nucleophilic Fluoride Sources. *Chem. Rev.* **2015**, *115*, 566.
- (8) Zhu, Q.; Gentry, E. C.; Knowles, R. R. Catalytic Carbocation Generation Enabled by the Mesolytic Cleavage of Alkoxyamine Radical Cations. *Angew. Chem., Int. Ed.* **2016**, *55*, 9969.
- (9) Xiang, J.; Shang, M.; Kawamata, Y.; Lundberg, H.; Reisberg, S. H.; Chen, M.; Mykhailiuk, P.; Beutner, G.; Collins, M. R.; Davies, A.; Del Bel, M.; Gallego, G. M.; Spangler, J. E.; Starr, J.; Yang, S.; Blackmond, D. G.; Baran, P. S. Hindered Dialkyl Ether Synthesis with Electrogenated Carbocations. *Nature* **2019**, *573*, 398.
- (10) Shibutani, S.; Kodo, T.; Takeda, M.; Nagao, K.; Tokunaga, N.; Sasaki, Y.; Ohmiya, H. Organophotoredox-Catalyzed Decarboxylative C(sp<sup>3</sup>)-O Bond Formation. *J. Am. Chem. Soc.* **2020**, *142*, 1211.
- (11) Atom transfer radical addition (ATRA) commonly invokes radical oxidation by a photocatalyst: (a) Wallentin, C. J.; Nguyen, J. D.; Finkbeiner, P.; Stephenson, C. R. J. Visible Light-Mediated Atom Transfer Radical Addition via Oxidative and Reductive Quenching of Photocatalysts. *J. Am. Chem. Soc.* **2012**, *134*, 8875. (b) Tlahuext-Aca, A.; Garza-Sanchez, R. A.; Glorius, F. Multicomponent Oxyalkylation of Styrenes Enabled by Hydrogen-Bond-Assisted Photoinduced Electron Transfer. *Angew. Chem., Int. Ed.* **2017**, *56* (13), 3708. For a review of ATRA: (c) Courant, T.; Masson, G. Recent Progress in Visible-Light Photoredox-Catalyzed Intermolecular 1,2-Difunctionalization of Double Bonds via an ATRA-Type Mechanism. *J. Org. Chem.* **2016**, *81*, 6945. ATRA-based fluorination: (d) Deng, W.; Feng, W.; Li, Y.; Bao, H. Merging Visible-Light Photocatalysis and Transition-Metal Catalysis in Three-Component Alkyl-Fluorination of Olefins with a Fluoride Ion. *Org. Lett.* **2018**, *20*, 2425. For radical-polar crossover to form anionic species: (e) Donabauer, K.; Maity, M.; Berger, A. L.; Huff, G. S.; Crespi, S.; König, B. Photocatalytic Carbanion Generation-Benzoylation of Aliphatic Aldehydes to Secondary Alcohols. *Chem. Sci.* **2019**, *10* (19), 5162.
- (12) For a recent review of transformations using redox active esters, see: Murarka, S. N-(Acyloxy)phthalimides as Redox-Active Esters in Cross-Coupling Reactions. *Adv. Synth. Catal.* **2018**, *360*, 1735.
- (13) (a) Wayner, D. D. M.; Houmam, A. Redox Properties of Free Radicals. *Acta Chem. Scand.* **1998**, *52*, 377. (b) Wayner, D. D. M.; McPhee, D. J.; Griller, D. Oxidation and Reduction Potentials of Transient Free Radicals. *J. Am. Chem. Soc.* **1988**, *110*, 132.
- (14) Anslyn, E.; Dougherty, D. A. *Modern physical organic chemistry*; University Science Books: 2006.
- (15) Silver-mediated decarboxylative fluorination with Selectfluor<sup>®</sup>: (a) Yin, F.; Wang, Z.; Li, Z.; Li, C. Silver-Catalyzed Decarboxylative Fluorination of Aliphatic Carboxylic Acids in Aqueous Solution. *J. Am. Chem. Soc.* **2012**, *134*, 10401. Photomediated decarboxylative fluorination with Selectfluor and NFSI: (b) Leung, J. C. T.; Chatalova-Sazepin, C.; West, J. G.; Rueda-Becerril, M.; Paquin, J. F.; Sammis, G. M. Photo-fluorodecarboxylation of 2-Aryloxy and 2-Aryl Carboxylic Acids. *Angew. Chem., Int. Ed.* **2012**, *51*, 10804. Photoredox-mediated decarboxylative fluorination: (c) Rueda-Becerril, M.; Mahé, O.; Drouin, M.; Majewski, M. B.; West, J. G.; Wolf, M. O.; Sammis, G. M.; Paquin, J. F. Direct C-F Bond Formation Using Photoredox Catalysis. *J. Am. Chem. Soc.* **2014**, *136*, 2637. (d) Leung, J. C. T.; Sammis, G. M. Radical Decarboxylative Fluorination of Aryloxyacetic Acids Using N-Fluorobenzenesulfonimide and a Photosensitizer. *Eur. J. Org. Chem.* **2015**, 2015, 2197. (e) Ventre, S.; Petronijevic, F. R.; Macmillan, D. W. C. Decarboxylative Fluorination of Aliphatic Carboxylic Acids via Photoredox Catalysis. *J. Am. Chem. Soc.* **2015**, *137*, 5654. For



- multi-fluorinated motifs, see: (f) Chatalova-Sazepin, C.; Binayeva, M.; Epifanov, M.; Zhang, W.; Foth, P.; Amador, C.; Jagdeo, M.; Boswell, B. R.; Sammis, G. M. Xenon Difluoride Mediated Fluorodecarboxylations for the Syntheses of Di- and Trifluoromethoxyarene. *Org. Lett.* **2016**, *18*, 4570. (g) Mizuta, S.; Stenhagen, I. S. R.; O'Duill, M.; Wolstenhulme, J.; Kirjavainen, A. K.; Forsback, S. J.; Tredwell, M.; Sandford, G.; Moore, P. R.; Huiban, M.; Luthra, S. K.; Passchier, J.; Solin, O.; Gouverneur, V. Catalytic Decarboxylative Fluorination for the Synthesis of Tri- and Difluoromethyl Arenes. *Org. Lett.* **2013**, *15*, 2648. (h) Wang, Z.; Guo, C. Y.; Yang, C.; Chen, J. P. Ag-Catalyzed Chemoselective Decarboxylative Mono- and gem- Difluorination of Malonic Acid Derivatives. *J. Am. Chem. Soc.* **2019**, *141*, 5617.
- (16) (a) Huang, X.; Liu, W.; Hooker, J. M.; Groves, J. T. Targeted Fluorination with the Fluoride Ion by Manganese-Catalyzed Decarboxylation. *Angew. Chem., Int. Ed.* **2015**, *54*, 5241. For a recent application of the methodology to the synthesis of multi-fluorinated motifs: (b) Sap, J. B. I.; Wilson, T. C.; Kee, C. W.; Straathof, N. J. W.; Ende, C. W. am; Mukherjee, P.; Zhang, L.; Genicot, C.; Gouverneur, V. Synthesis of  $^{18}\text{F}$ -Difluoromethylarenes Using Aryl Boronic Acids, Ethyl Bromofluoroacetate and  $[\text{F}^-]$ -Fluoride. *Chem. Sci.* **2019**, *10*, 3237.
- (17) Gray, E. E.; Nielsen, M. K.; Choquette, K. A.; Kalow, J. A.; Graham, T. J. A.; Doyle, A. G. Nucleophilic (Radio)Fluorination of  $\alpha$ -Diazocarbonyl Compounds Enabled by Copper-Catalyzed H-F Insertion. *J. Am. Chem. Soc.* **2016**, *138*, 10802.
- (18) (a) Ametamey, S. M.; Honer, M.; Schubiger, P. A. Molecular Imaging with PET. *Chem. Rev.* **2008**, *108*, 1501. (b) Cole, E.; Stewart, M.; Littich, R.; Hoareau, R.; Scott, P. Radiosyntheses Using Fluorine-18: The Art and Science of Late Stage Fluorination. *Curr. Top. Med. Chem.* **2014**, *14*, 875–900. (c) Cai, L.; Lu, S.; Pike, V. W. Chemistry with  $[\text{F}^-]$  Fluoride Ion. *Eur. J. Org. Chem.* **2008**, *2008*, 2853–2873. (d) Jacobson, O.; Kiesewetter, D. O.; Chen, X. Fluorine-18 Radiochemistry, Labeling Strategies and Synthetic Routes. *Bioconjugate Chem.* **2015**, *26*, 1–18. (e) Deng, X.; Rong, J.; Wang, L.; Vasdev, N.; Zhang, L.; Josephson, L.; Liang, S. H. Chemistry for Positron Emission Tomography: Recent Advances in  $^{11}\text{C}$ -,  $^{18}\text{F}$ -,  $^{13}\text{N}$ -, and  $^{15}\text{O}$ -Labeling Reactions. *Angew. Chem., Int. Ed.* **2019**, *58*, 2580–2605. (f) Liang, S. H.; Vasdev, N. Total Radiosynthesis: Thinking Outside 'the Box'. *Aust. J. Chem.* **2015**, *68*, 1319–1328.
- (19) (a) Teegardin, K.; Day, J. I.; Chan, J.; Weaver, J. Advances in Photocatalysis: A Microreview of Visible Light Mediated Ruthenium and Iridium Catalyzed Organic Transformations. *Org. Process Res. Dev.* **2016**, *20*, 1156. (b) Speckmeier, E.; Fischer, T. G.; Zeitler, K. A Toolbox Approach to Construct Broadly Applicable Metal-Free Catalysts for Photoredox Chemistry: Deliberate Tuning of Redox Potentials and Importance of Halogens in Donor-Acceptor Cyanoarenes. *J. Am. Chem. Soc.* **2018**, *140* (45), 15353.
- (20) Okada, K.; Okamoto, K.; Morita, N.; Okubo, K.; Oda, M. Photosensitized Decarboxylative Michael Addition through N-(Acyloxy)Phthalimides via an Electron-Transfer Mechanism. *J. Am. Chem. Soc.* **1991**, *113*, 9401.
- (21) Boutureira, O.; Bernardes, G. J. L. Advances in Chemical Protein Modification. *Chem. Rev.* **2015**, *115*, 2174.
- (22) Shimizu, M.; Hiyama, T. Modern Synthetic Methods for Fluorine-Substituted Target Molecules. *Angew. Chem., Int. Ed.* **2005**, *44*, 214.
- (23) Zafrani, Y.; Yeffet, D.; Sod-Moriah, G.; Berliner, A.; Amir, D.; Marciano, D.; Gershonov, E.; Saphier, S. Difluoromethyl Bioisostere: Examining the "Lipophilic Hydrogen Bond Donor" Concept. *J. Med. Chem.* **2017**, *60* (2), 797–804.
- (24) Nielsen, M. K.; Ahneman, D. T.; Riera, O.; Doyle, A. G. Deoxyfluorination with Sulfonyl Fluorides: Navigating Reaction Space with Machine Learning. *J. Am. Chem. Soc.* **2018**, *140* (15), 5004.
- (25) González-Esguevillas, M.; Miró, J.; Jeffrey, J. L.; MacMillan, D. W. C. Photoredox-catalyzed Deoxyfluorination of Activated Alcohols with Selectfluor. *Tetrahedron* **2019**, *75*, 4222.
- (26) (a) Lackner, G. L.; Quasdorf, K. W.; Overman, L. E. Direct Construction of Quaternary Carbons from Tertiary Alcohols via Photoredox-Catalyzed Fragmentation of *tert*-Alkyl N-Phthalimidoyl Oxalates. *J. Am. Chem. Soc.* **2013**, *135*, 15342. (b) Lackner, G. L.; Quasdorf, K. W.; Pratsch, G.; Overman, L. E. Fragment Coupling and the Construction of Quaternary Carbons Using Tertiary Radicals Generated from *tert*-Alkyl N-Phthalimidoyl Oxalates by Visible-Light Photocatalysis. *J. Org. Chem.* **2015**, *80*, 6012.
- (27) See Supporting Information for additional details and discussion.
- (28) Cismesia, M. A.; Yoon, T. P. Characterizing Chain Processes in Visible Light Photoredox Catalysis. *Chem. Sci.* **2015**, *6*, 5426.
- (29) An acidic additive was required in order to observe reactivity with these alternative nucleophiles. No conversion was observed without  $\text{Et}_3\text{N}\cdot 3\text{HF}$  except in the case of sufficiently acidic HFIP.
- (30) (a) Zlatopolskiy, B. D.; Zischler, J.; Krapf, P.; Zarrad, F.; Urusova, E. A.; Kordys, E.; Endepols, H.; Neumaier, B. Copper-Mediated Aromatic Radiofluorination Revisited: Efficient Production of PET Tracers on a Preparative Scale. *Chem. - Eur. J.* **2015**, *21*, 5972. (b) Jacobson, O.; Kiesewetter, D. O.; Chen, X. Fluorine-18 Radiochemistry, Labeling Strategies and Synthetic Routes. *Bioconjugate Chem.* **2015**, *26*, 1.
- (31) D'Angelo, K. A.; Taylor, M. S. Borinic Acid Catalyzed Stereo- and Regioselective Couplings of Glycosyl Methanesulfonates. *J. Am. Chem. Soc.* **2016**, *138*, 11058.