

Phosphine/Photoredox Catalyzed Anti-Markovnikov Hydroamination of Olefins with Primary Sulfonamides via α -Scission from Phosphoranyl Radicals

Alex J. Chinn,[#] Kassandra Sedillo,[#] and Abigail G. Doyle*Cite This: <https://doi.org/10.1021/jacs.1c09484>

Read Online

ACCESS |




Metrics & More



Article Recommendations



Supporting Information



ABSTRACT: New strategies to access radicals from common feedstock chemicals hold the potential to broadly impact synthetic chemistry. We report a dual phosphine and photoredox catalytic system that enables direct formation of sulfonamidyl radicals from primary sulfonamides. Mechanistic investigations support that the N-centered radical is generated via α -scission of the P–N bond of a phosphoranyl radical intermediate, formed by sulfonamide nucleophilic addition to a phosphine radical cation. As compared to the recently well-explored β -scission chemistry of phosphoranyl radicals, this strategy is applicable to activation of N-based nucleophiles and is catalytic in phosphine. We highlight application of this activation strategy to an intermolecular anti-Markovnikov hydroamination of unactivated olefins with primary sulfonamides. A range of structurally diverse secondary sulfonamides can be prepared in good to excellent yields under mild conditions.

INTRODUCTION

Photoredox catalysis and electrochemistry have enabled the development of valuable synthetic methods by offering mild, selective, and practical mechanisms for electron transfer.^{1,2} While many common functional groups can be easily oxidized or reduced to afford reactive radical intermediates, many others remain inaccessible or present chemoselectivity challenges due to prohibitively high oxidation/reduction potentials.³ A classic strategy to circumvent these challenges is to append a sacrificial redox auxiliary to the functional group to alter its redox potential, albeit this strategy introduces additional synthetic manipulations as well as accompanying waste and scope limitations.⁴ Recently, alternative approaches, such as *in situ* activation,⁵ proton-coupled electron transfer (PCET),⁶ molecular electrocatalysis,^{2,7} electro-photochemistry,⁸ and multiphoton excitation, have been explored to enable the generation of valuable radical classes otherwise challenging to access directly from feedstock chemicals via electron transfer.⁹

In this context, our group and others were drawn to phosphoranyl radicals to effect homolytic cleavage of otherwise redox inaccessible X–Y bonds via photoredox catalysis.^{10,11} Seminal publications from Benetrupe and Roberts demon-

strated that these intermediates fragment through either an α - or β -scission pathway to produce a new radical and a resulting P(III) or P(V) species, respectively (Figure 1A).¹² Historically, phosphoranyl radicals were accessed by direct radical addition to a P(III) species, and the harsh conditions required for radical generation limited widespread synthetic applications.¹³ More recently, photoredox catalysis has provided a platform to generate phosphoranyl radicals under mild conditions via either radical addition to the P(III) species or single-electron oxidation of the P(III) species followed by nucleophilic addition (Figure 1B).¹⁴ Thus, this activation strategy has seen recent widespread application for deoxygenation and desulfurization reactions of various common organic functional groups.¹⁵ As an example, our group recently described the direct deoxygenation of benzylic alcohols and carboxylic acids

Received: September 7, 2021






ACS Publications

© XXXX American Chemical Society

A

<https://doi.org/10.1021/jacs.1c09484>
J. Am. Chem. Soc. XXXX, XXX, XXX–XXX

A. Fragmentation pathways of phosphoranyl radicals

B. Prior art: photoredox / phosphine mediated R-X activation via β -scissionC. Proposal: photoredox / phosphine catalyzed X-H activation via α -scission

D. This work: Intermolecular hydroamination of unactivated olefins



Figure 1. Synthesis and reactivity of phosphoranyl radicals.

and the hydroacylation of olefins with carboxylic acids using a phosphine in conjunction with visible light photoredox catalysis.¹⁰

While β -scission methods have proven to be effective in overcoming voltage-gated restrictions, the process is thermodynamically driven by formation of the stoichiometric phosphine oxide byproduct, which has largely limited its utility to oxygen-based nucleophiles. We questioned whether taking advantage of the synthetically unexplored α -scission fragmentation pathway, wherein scission is instead kinetically favored due to an unpaired electron occupying a P–X antibonding orbital, could lead to new opportunities for synthesis and facilitate a process catalytic in phosphine (Figure 1C). Prior studies have shown that α -scission occurs site selectively from the apical position of a phosphoranyl radical and that electronegative functional groups have a preference for the apical site, suggesting that electronegative ligands could selectively undergo α -scission.^{12d,e,16} As such, we envisioned a process wherein a trapped nucleophile would preferentially undergo α -scission, without displacing any P(III) substituents already present, enabling the generation of radical species with only catalytic amounts of phosphine. While the generation of P=O/S bonds via β -scission could potentially compete with

α -scission for many O- and S-nucleophiles, we hypothesized that a catalytic α -scission strategy could be applied to N-based nucleophiles, for which no examples of β -scission have been reported. With these criteria in mind, we were drawn to study N–H bond activation of primary sulfonamides, which possess both high N–H bond strength and oxidation potential (BDFE ~ 105 kcal/mol, $E_{1/2} = +2.6$ V versus SCE in MeCN). Here we demonstrate realization of this goal in the context of an anti-Markovnikov olefin hydroamination with primary sulfonamides catalyzed by both tricyclohexylphosphine (PCy₃) and a visible light photoredox catalyst (Figure 1D).


Intermolecular hydroamination reactions between primary sulfonamides and olefins are an attractive method for the direct and atom-economical synthesis of secondary sulfonamides, a prevalent structural feature in bioactive molecules.¹⁷ Whereas Markovnikov-selective approaches have significant precedent,¹⁸ only recently have methods accessing anti-Markovnikov selectivity been described. In particular, Nicewicz and co-workers described a photoredox-catalyzed anti-Markovnikov hydroamination that proceeds via alkene radical cation intermediates.¹⁹ Whereas this strategy is applicable to trisubstituted aliphatic alkenes and substituted styrenes, Knowles and co-workers showed that anti-Markovnikov hydroamination of unactivated olefins was possible via PCET activation of the N–H bonds to produce a sulfonamidyl radical.²⁰ While highly enabling, the intermolecular reaction has not been extensively demonstrated beyond electron-rich *p*-methoxybenzenesulfonamide. Thus, identification of new catalytic strategies for N–H bond activation could offer synthetic advances in catalytic olefin hydroamination.^{21,22}

RESULTS AND DISCUSSION

We envisioned a catalytic cycle beginning with excitation of the photocatalyst, followed by single electron oxidation of a phosphine catalyst A to the corresponding radical cation B (Figure 2A). Nucleophilic trapping of B with sulfonamide D would form phosphoranyl radical intermediate C. α -Scission of the P–N bond in C would liberate an N-centered radical E and regenerate the phosphine catalyst A. Reaction of the N-centered radical with the olefin partner would form intermediate F, which, upon hydrogen atom transfer (HAT) with catalytic thiol H, should generate the desired product G. Single electron transfer (SET) between the thiyl radical I and the reduced photocatalyst, followed by proton transfer (PT), would regenerate the photocatalyst as well as the HAT catalyst, completing the cycle.

Notably, success in the proposed scheme would require that reaction conditions are carefully tuned to accommodate the promiscuous reactivity of phosphine radical cation B and phosphoranyl radical C (Figure 2B). While nucleophilic trapping of a sulfonamide by B would progress the reaction (i), the presence of other nucleophilic species, such as excess P(III) A and thiol H, could lead to unproductive phosphoranyl radical formation (ii) and subsequent degradation.²³ Phosphine radical cations have also been shown to react with olefins, which could then form the corresponding phosphonium species upon HAT (iii).²⁴ From the phosphoranyl radical C, undesired decomposition such as overoxidation to form imidophosphorane K is possible (iv).²⁵ Moreover, while sulfonamide α -scission should be kinetically preferred (v), undesired P–R bond α -scission (vi) could give rise to aminophosphorane L. These pathways are likely to dominate if P–N bond α -scission is inefficient or if the N-centered

A. Proposed catalytic cycle



B. Phosphine speciation pathways





Figure 2. Proposed catalytic cycle and alternative speciation pathways.

radical is not rapidly trapped by the olefin, leading to reformation of the phosphoranyl radical.

With these considerations in mind, we embarked on initial reaction discovery efforts using the hydroamination of electronically unactivated 1-hexene with *p*-*tert*-butylbenzenesulfonamide (**1**) as a model system (Table 1). On the basis of our prior work,^{10a} we elected to evaluate [Ir(dF(Me)ppy)₂-PF₆] (3) [dF(Me)ppy = 2-(2,4-difluorophenyl)-5-methylpyridine; dtbbpy = 4,4'-di-*tert*-butyl-2,2'-bipyridine] as a photocatalyst and lutidine as a base to facilitate proton transfer. Sterically hindered 2,4,6-triisopropylbenzenethiol (TRIP-SH) was selected as an HAT catalyst to sterically disfavor nucleophilic addition of the thiol to the phosphine radical cation. To our delight, we found that triphenylphosphine catalyzed the hydroamination in PhCF₃ as solvent, albeit in low yield (12%, entry 1). Electron-rich P(*p*-MeOC₆H₄)₃ showed no improvement in yield (12%), and electron-deficient P(*p*-CF₃C₆H₄)₃ afforded no product (entries 2 and 3). We were excited to observe that trialkylphosphines exhibited improved levels of reactivity (entries 4–7): PCy₃, which is commercially available and relatively inexpensive, provided the highest yield at 46%.²⁶


Next, we evaluated other reaction dimensions using PCy₃ as the catalyst. Hydroamination proceeded in the absence of lutidine (entry 8), likely because PCy₃ itself has comparable basicity (pK_a ~ 15 in MeCN) and can facilitate proton transfer with TRIP thiol.²⁷ Inclusion of a stronger base, such as the

Table 1. Optimization Studies



entry ^a	phosphine (mol%)	base	photocatalyst	yield (%) ^b
1	PPh ₃ (10 mol%)	Lutidine	3	12
2	P(4-MeOC ₆ H ₄) ₃ (10 mol%)	Lutidine	3	12
3	P(4-CF ₃ C ₆ H ₄) ₃ (10 mol%)	Lutidine	3	0
4	PMe ₃ (10 mol%)	Lutidine	3	34
5	PEt ₃ (10 mol%)	Lutidine	3	41
6	P(<i>t</i> -Bu) ₃ (10 mol%)	Lutidine	3	37
7	PCy ₃ (10 mol%)	Lutidine	3	46
8	PCy ₃ (10 mol%)	none	3	46
9	PCy ₃ (10 mol%)	NBu ₄ OP(O)(OBu) ₂	3	32
10	PCy ₃ (5 mol%)	none	3	60
11	PCy ₃ (2.5 mol%)	none	3	63
12 ^c	PCy ₃ (2.5 mol%)	none	3	87
13	PCy ₃ (2.5 mol%)	none	4	0
14	PCy ₃ (2.5 mol%)	none	5	49
15	PCy ₃ (2.5 mol%)	none	6	10
16 ^d	PCy ₃ (2.5 mol%)	none	3	79

entry ^{a,c}	deviation from best conditions (entry 10)	yield (%) ^b
17	no Phosphine	0
18	no Photocatalyst	0
19	no TRIP-SH	3
20	no Light	0



3: R¹ = *t*-Bu, R² = Me, R³ = F

Ir^{III}/*Ir^{III} = +0.97 V vs SCE

4: R¹ = *t*-Bu, R² = H, R³ = H

Ir^{III}/*Ir^{III} = +0.66 V vs SCE

5: R¹ = *t*-Bu, R² = CF₃, R³ = F

Ir^{III}/*Ir^{III} = +1.21 V vs SCE

6: R¹ = CF₃, R² = CF₃, R³ = F

Ir^{III}/*Ir^{III} = +1.65 V vs SCE

^aReactions were performed on a 0.1 mmol scale with 1.0 equiv of sulfonamide **1** and 1.0 equiv of 1-hexene by using blue LEDs as a light source. ^bYield determined by ¹H NMR by comparison to 1,4-bis(trimethylsilyl)benzene as an internal standard and are reported as an average of two runs. ^cReaction was run with 2.0 equiv of olefin. ^dReaction was conducted on a 0.5 mmol scale by using a Kessil H150 lamp for 24 h.


monophosphate base employed by Knowles and co-workers in their PCET hydroamination, led to decreased reaction yield, potentially due to promotion of undesired nucleophilic addition into the phosphine radical cation (ii in Figure 2B). Further experimentation revealed that the phosphine catalyst loading has a significant impact on reactivity, with lower loadings of phosphine showing an increase in yield (entries 8, 10, and 11). We attribute this observation to the fact that higher concentrations of phosphine could favor addition of the N-centered radical to PCy₃ to reform the phosphoranyl radical according to LeChatelier's principle (Figure 2B, reverse of vi), which could ultimately favor catalyst decomposition according

to pathways iv and vi. We hypothesized that an increase in olefin stoichiometry could kinetically trap the N-centered radical, potentially outcompeting the decomposition pathways. Gratifyingly, reactions performed with 2 equiv of olefin delivered product **2** in 87% yield (entry 12).²⁸ Interestingly, despite the use of excess olefin, only monofunctionalized sulfonamide was observed, likely because the product is too sterically hindered to undergo nucleophilic addition to the phosphine radical cation. Upon evaluating a selection of Ir photocatalysts, we found that those with excited state oxidation potentials below that of PCy₃ were completely ineffective (entry 13). Catalysts with higher oxidation potentials also led to decreased yields likely due to over oxidation of the phosphoranyl radical to form the imidophosphorane **K** (entries 14 and 15; *vide infra* for full discussion of the catalytic relevance of **K**). Finally, reactions run on a 0.5 mmol scale were conducted by using Kessil lamps (34 W) rather than blue LEDs (12 W), resulting in a notable increase in yield (entry 16). Control experiments revealed that each component was required to achieve high levels of reactivity (entries 17–20).


Although our observations during reaction optimization were consistent with a phosphoranyl radical mechanism, alternative reaction pathways could lead to product formation. Direct oxidation/deprotonation could instead generate a sulfonamidyl radical, or a multisite PCET pathway (Figure 3A), as demonstrated by Knowles and co-workers, could be operative. Alternatively, it is possible that C–N bond formation occurs through a reductive elimination mechanism from a P(V) intermediate analogous to work from McNally and co-workers in C–C, C–O, and C–N bond formation (Figure 3B).³⁵ Such a species could arise from either nucleophilic addition of a sulfonamide to phosphonium **J** or a two-step olefin addition/HAT process with a phosphoranyl radical.

The high redox potential and pK_a of sulfonamides ($E_{1/2} = +2.6$ V versus SCE in MeCN, pK_a ~ 27 in MeCN)²⁰ eliminate the possibility of a direct oxidation/deprotonation mechanism. Instead, if multisite PCET were operative, the basic phosphine and photocatalyst would work in concert to perform a net HAT (Figure 3A).²⁹ We reasoned that Stern–Volmer quenching studies would be effective in discriminating between this pathway and a phosphoranyl radical mechanism.³⁰ If a PCET pathway is operative, then a combination of sulfonamide and phosphine would be more effective at quenching an excited photocatalyst than either component on its own. Instead, we found that PCy₃ was more effective at quenching the excited photocatalyst (Figure 3C, red circles, $K_{SV} = 1617$) than the combination of the phosphine and sulfonamide, which produced a smaller slope (green squares, $K_{SV} = 1486$) (Figure 3C, left). Studies conducted wherein sulfonamide concentration was increased in the presence of constant phosphine showed a slightly negative slope ($K_{SV} = -0.02$) (Figure 3C, right). These experiments are inconsistent with a PCET mechanism and suggest instead that hydrogen bonding between the phosphine and sulfonamide removes electron density from the phosphine, making it a less effective quenching species.³¹ We also assessed the effective BDFE of the oxidant/base pair, as demonstrated by Mayer and co-workers.³² A value of ~90 kcal/mol is found for PCy₃ and [Ir(dF(Me)ppy)₂(dtbbpy)]PF₆, whereas the calculated BDFE of a primary sulfonamide is ~105 kcal/mol, making a PCET mechanism thermodynamically unlikely.^{20,33} Instead, the rapid oxidation of phosphine in these studies indicates that a


A. Alternative mechanism: N–H activation via proton-coupled electron transfer




B. Alternative mechanism: C–N bond formation from P(V) intermediate



C. Stern–Volmer luminescence quenching experiments



D. Catalytic performance of isolated P-derived byproducts



E. Evaluation of P(V) mechanism with mixed *n*-alkyl ligands




Figure 3. Mechanistic studies. ^aStandard conditions: as in entry 16, Table 1 without PCy₃.

phosphine radical cation is readily formed in the reaction mixture, which should facilitate phosphoranyl radical formation.

Although preliminary efforts to directly observe the phosphoranyl radical intermediate were unsuccessful, we were able to detect via ³¹P NMR aminophosphine **L-1** and imidophosphorane **K-1**, which likely arise via the pathways outlined in Figure 2B (Figure 3D) (see section 4.1 of the Supporting Information for full details).³⁴ While these species provide indirect evidence for the intermediacy of a phosphoranyl radical intermediate, we also recognized that they could conceivably be catalytically relevant either as catalysts themselves or as bases in multisite PCET. However,

subjecting independently synthesized samples of **L-1** and **K-1** to the reaction conditions in the absence of PCy_3 revealed that neither species was catalytically viable (Figure 3D).

While multiple examples of C–C bond formation via P(V) reductive elimination have been reported,³⁶ no examples exist for C(sp³)–N bond formation. Nevertheless, we sought to experimentally evaluate the feasibility of the reductive C–N bond formation pathway in Figure 3B. First, we subjected independently prepared phosphonium **J-1** to the catalytic reaction conditions in the absence of PCy_3 (Figure 3D). Hydroamination was not observed, suggesting that the nucleophilic addition pathway to generate the P(V) intermediate is not viable, although counterion effects may influence reactivity. While the olefin addition/HAT pathway is still possible, to the best of our knowledge, there are no reports wherein phosphoranyl radicals have been trapped by olefins, making this pathway unlikely. To investigate the proposed P(V) mechanism further, triethylphosphine was used as the catalyst under standard reaction conditions (Figure 3E). This catalyst would afford P(V) intermediate **8**, from which a mixture of ethylated and *n*-hexylated secondary sulfonamides (**7** and **2**) would be expected since there should be little to no difference in reductive elimination among these two primary alkyl substituents. However, formation of desired product **2** occurred in 50% yield while the ethylated product **7** was not observed by ¹H NMR nor HRMS (see section 4.5 of the Supporting Information for full details). Together these studies provide strong evidence that the proposed α -scission mechanism is more likely than either a PCET or a P(V) reductive elimination and that PCy_3 is likely the active form of the catalyst.

We next examined a variety of olefins under our optimized reaction conditions. Acyclic, cyclic, and bicyclic olefins underwent hydroamination to afford the desired products in high yield (Figure 4). Terminal monosubstituted, 1,1-disubstituted, and 1,2-disubstituted olefins were also competent reaction partners (**2**, **9**–**13**). In the case of *cis*- and *trans*-4-octene, we found that alkene configuration does not affect reactivity (**14**, **15**). Trisubstituted and tetrasubstituted olefins were also competent substrates (**16**, **17**), albeit hydroamination proceeded in lower yields likely because sulfonamidyl radical trapping is not as efficient with these more hindered olefins, leading to reformation of the phosphoranyl radical and subsequent decomposition pathways (Figure 2B). In general, the electrophilic sulfonamidyl radical undergoes hydroamination in good yield with electron-rich olefins, such as an enol ether (**18**) and vinylpyrrolidinone (**19**).³⁷ Only modest reactivity is observed with styrene (**20**) and allylbenzene (**21**).

Gratifyingly, the standard reaction conditions tolerate a wide range of functional groups, including carbamates (**22**), primary alkyl chlorides (**23**), unprotected primary alcohols (**24**), and methyl esters (**25**). The success of **24** is particularly encouraging as no competitive β -scission was detected despite the presence of a nucleophilic alcohol.

Turning to the sulfonamide scope, we found that hydroamination of cyclohexene with benzenesulfonamide under the standard reaction conditions delivered only 54% of the desired product **26** (Figure 5). We hypothesized that electron-neutral and deficient aryl sulfonamides and alkyl sulfonamides might undergo α -scission less efficiently due to the relative instability of the resulting N-centered radical. As such, we reasoned that reactivity could be improved by employing an excess of olefin




Figure 4. Olefin scope. ^aReactions performed on a 0.5 mmol scale with 1.0 equiv of sulfonamide **1** (Ar = *p*-*tert*-butylbenzene) and 1.0 equiv of olefin. Isolated yield, reported as an average of two runs. ^b1:2 *cis/trans* mixture of diastereomers. ^cReaction was conducted with 3.0 equiv of olefin. ^d¹H NMR yield determined by comparison to an internal standard.

to rapidly trap the sulfonamidyl radical and prevent decomposition of the catalyst. Because we previously observed that the method is selective for monoalkylation, we anticipated that overalkylation would not be an issue. While employment of an excess of olefin is not ideal, sulfonamides are often considered the more valuable component in drug discovery.^{17c,d,38}

Gratifyingly, upon employing 5.0 equiv of cyclohexene, we were able to isolate the secondary sulfonamides derived from electron-rich and neutral substrates (**11**, **26**–**28**) in 74–85% yield. However, an electron-deficient aryl sulfonamide delivered low yield (**29**), likely resulting from poor nucleophilicity of the substrate which limits its ability to trap the phosphine radical cation. Using these conditions, we were pleased to see a variety of medically relevant motifs undergo hydroamination. For example, *o*-chlorobenzenesulfonamides (**30**), a structure prevalent in 12 approved drugs,³⁹ proved to be a competent coupling partner (70% yield). Additionally, thiophene-2-sulfonamide, a scaffold in the approved drugs dorzolamide and brinzolamide, and *N,N*-dimethylsulfamoylamine, present in the approved drug beclabuvir, gave products **31** and **32**, respectively. The catalytic reaction can be extended to aliphatic sulfonamides as well. Cyclopropyl and methyl sulfonamide underwent olefin hydroamination to afford **33** and **34** in 68% and 89% yield, respectively.




Figure 5. Sulfonamide scope. ^aReactions were performed on a 0.5 mmol scale. Isolated yield, reported as an average of two runs. ^b¹H NMR yield determined by comparison to 1,4-bis(trimethylsilyl)-benzene as an internal standard and reported as an average of two runs. ^cReaction was conducted with 1.0 equiv of olefin.

CONCLUSION

In summary, we have developed a catalytic strategy for N–H activation of sulfonamides via phosphine and photoredox catalysis. This strategy was explored in an intermolecular anti-Markovnikov hydroamination of unactivated olefins with sulfonamides. Mechanistic studies suggest that sulfonamidyl radical generation proceeds via α -scission from a phosphoranyl radical intermediate in a polar-radical crossover pathway. We anticipate that with improved understanding of the reactivity and selectivity of phosphoranyl radicals it will be possible to further expand the diversity of nucleophiles and reaction classes capable of interfacing with phosphine/photoredox catalysis and apply this strategy in new synthetic contexts.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/jacs.1c09484>.

Experimental procedures, experimental data, and characterization and spectral data for new compounds (PDF)

AUTHOR INFORMATION

Corresponding Author

Abigail G. Doyle – Department of Chemistry, Princeton University, Princeton, New Jersey 08544, United States; Department of Chemistry and Biochemistry, University of California, Los Angeles, Los Angeles, California 90095, United States; orcid.org/0000-0002-6641-0833; Email: agdoyle@chem.ucla.edu

Authors

Alex J. Chinn – Department of Chemistry, Princeton University, Princeton, New Jersey 08544, United States

Kassandra Sedillo – Department of Chemistry, Princeton University, Princeton, New Jersey 08544, United States

Complete contact information is available at: <https://pubs.acs.org/doi/10.1021/jacs.1c09484>

Author Contributions

#A.J.C. and K.F.S. contributed equally to this work.

Funding

Financial support for this project was provided by the BioLEC, an Energy Frontier Research Center funded by the U.S. Department of Energy, Office of Science, Office of Basic Energy Sciences, under Award DE-SC0019370.

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

We thank Dr. John Eng for assistance with analytical equipment and Dr. Jesus I. Martinez-Alvarado for helpful discussions.

REFERENCES

- (1) (a) Shaw, M. H.; Twilton, J.; MacMillan, D. W. Photoredox Catalysis in Organic Chemistry. *J. Org. Chem.* **2016**, *81*, 6898–6926. (b) Romero, N. A.; Nicewicz, D. A. Organic Photoredox Catalysis. *Chem. Rev.* **2016**, *116*, 10075–10166. (c) Crespi, S.; Fagnoni, M. Generation of Alkyl Radicals: From the Tyranny of Tin to the Photon Democracy. *Chem. Rev.* **2020**, *120*, 9790–9833. (d) Narayanam, J. M. R.; Stephenson, C. R. J. Visible Light Photoredox Catalysis: Applications in Organic Synthesis. *Chem. Soc. Rev.* **2011**, *40*, 102–113. (e) Goddard, J.-P.; Ollivier, C.; Fensterbank, L. Photoredox Catalysis for the Generation of Carbon Centered Radicals. *Acc. Chem. Res.* **2016**, *49*, 1924–1936.
- (2) (a) Yan, M.; Kawamata, Y.; Baran, P. S. Synthetic Organic Electrochemical Methods Since 2000: On the Verge of a Renaissance. *Chem. Rev.* **2017**, *117*, 13230–13319. (b) Zhu, C.; Ang, N. W. J.; Meyer, T. H.; Qiu, Y.; Ackermann, L. Organic Electrochemistry: Molecular Syntheses with Potential. *ACS Cent. Sci.* **2021**, *7*, 415–431. (c) Shi, S.-H.; Liang, Y.; Jiao, N. Electrochemical Oxidation Induced Selective C–C Bond Cleavage. *Chem. Rev.* **2021**, *121*, 485–505.
- (3) Roth, H. G.; Romero, N. A.; Nicewicz, D. A. Experimental and Calculated Electrochemical Potentials of Common Organic Molecules for Applications to Single-Electron Redox Chemistry. *Synlett* **2016**, *27*, 714–723.
- (4) (a) Guo, J.-J.; Hu, A.; Zuo, Z. Photocatalytic Alkoxy Radical-Mediated Transformations. *Tetrahedron Lett.* **2018**, *59*, 2103–2111. (b) Chen, J. R.; Hu, X. Q.; Lu, L. Q.; Xiao, W. J. Visible Light Photoredox-Controlled Reactions of N-Radicals and Radical Ions. *Chem. Soc. Rev.* **2016**, *45*, 2044–2056.
- (5) Wappes, E. A.; Nakafuku, K. M.; Nagib, D. A. Directed β C–H Amination of Alcohols via Radical Relay Chaperones. *J. Am. Chem. Soc.* **2017**, *139*, 10204–10207.
- (6) (a) Gentry, E. C.; Knowles, R. R. Synthetic Applications of Proton-Coupled Electron Transfer. *Acc. Chem. Res.* **2016**, *49*, 1546–1556. (b) Tsui, E.; Wang, H.; Knowles, R. R. Catalytic Generation of Alkoxy Radicals From Unfunctionalized Alcohols. *Chem. Sci.* **2020**, *11*, 11124–11141. (c) Thammavongsy, Z.; Mercer, I. P.; Yang, J. Y. Promoting Proton Coupled Electron Transfer in Redox Catalysts Through Molecular Design. *Chem. Commun.* **2019**, *55*, 10342–10358.
- (7) (a) Novaes, L. F. T.; Liu, J.; Shen, Y.; Lu, L.; Meinhardt, J. M.; Lin, S. Electrocatalysis as an Enabling Technology for Organic Synthesis. *Chem. Soc. Rev.* **2021**, *50*, 7941–8002. (b) Savéant, J.-M. Molecular Catalysis of Electrochemical Reactions. Mechanistic Aspects. *Chem. Rev.* **2008**, *108*, 2348–2378.
- (8) (a) Liu, J.; Lu, L.; Wood, D.; Lin, S. New Redox Strategies in Organic Synthesis by Means of Electrochemistry and Photochemistry. *ACS Cent. Sci.* **2020**, *6*, 1317–1340. (b) Yu, Y.; Guo, P.; Zhong, J.-S.;

- Yuan, Y.; Ye, K.-Y. Merging Photochemistry with Electrochemistry in Organic Synthesis. *Org. Chem. Front.* **2020**, *7*, 131–135. (c) Huang, H.; Strater, Z. M.; Rauch, M.; Shee, J.; Sisto, T. J.; Nuckolls, C.; Lambert, T. H. Electrophotocatalysis with a Trisaminocyclopropenium Radical Dication. *Angew. Chem., Int. Ed.* **2019**, *58*, 13318–13322. (d) Zhang, W.; Carpenter, K. L.; Lin, S. Electrochemistry Broadens the Scope of Flavin Photocatalysis: Photoelectrocatalytic Oxidation of Unactivated Alcohols. *Angew. Chem., Int. Ed.* **2020**, *59*, 409–417. (e) Wang, F.; Stahl, S. S. Merging Photochemistry with Electrochemistry: Functional-Group Tolerant Electrochemical Amination of C(sp³)–H Bonds. *Angew. Chem., Int. Ed.* **2019**, *58*, 6385–6390. (f) Shen, T.; Lambert, T. H. Electrophotocatalytic Diamination of Vicinal C–H Bonds. *Science* **2021**, *371*, 620–626. (g) Cowper, N. G. W.; Chernowsky, C. P.; Williams, O. P.; Wickens, Z. K. Potent Reductants via Electron-Primed Photoredox Catalysis: Unlocking Aryl Chlorides for Radical Coupling. *J. Am. Chem. Soc.* **2020**, *142*, 2093–2099.
- (9) (a) Glaser, F.; Kerzig, C.; Wenger, O. S. Multi-Photon Excitation in Photoredox Catalysis: Concepts, Applications, Methods. *Angew. Chem., Int. Ed.* **2020**, *59*, 10266–10284. (b) Ravetz, B. D.; Pun, A. B.; Churchill, E. M.; Congreve, D. N.; Rovis, T.; Campos, L. M. Photoredox Catalysis Using Infrared Light via Triplet Fusion Upconversion. *Nature* **2019**, *565*, 343–346.
- (10) (a) Stache, E. E.; Ertel, A. B.; Rovis, T.; Doyle, A. G. Generation of Phosphoranyl Radicals via Photoredox Catalysis Enables Voltage-Independent Activation of Strong C–O Bonds. *ACS Catal.* **2018**, *8*, 11134–11139. (b) Martinez Alvarado, J. I.; Ertel, A. B.; Stegner, A.; Stache, E. E.; Doyle, A. G. Direct Use of Carboxylic Acids in the Photocatalytic Hydroacylation of Styrenes To Generate Dialkyl Ketones. *Org. Lett.* **2019**, *21*, 9940–9944.
- (11) (a) Zhang, M.; Xie, J.; Zhu, C. A General Deoxygenation Approach for Synthesis of Ketones From Aromatic Carboxylic Acids and Alkenes. *Nat. Commun.* **2018**, *9*, 3517–3527. (b) Ruzi, R.; Ma, J.; Yuan, X.-A.; Wang, W.; Wang, S.; Zhang, M.; Dai, J.; Xie, J.; Zhu, C. Deoxygenative Arylation of Carboxylic Acids by Aryl Migration. *Chem. - Eur. J.* **2019**, *25*, 12724–12729. (c) Ning, Y.; Wang, S.; Li, M.; Han, J.; Zhu, C.; Xie, J. Site-Specific Umpolung Amidation of Carboxylic Acids, Via Triplet Synergistic Catalysis. *Nat. Commun.* **2021**, *12*, 4637–4636.
- (12) (a) Bentrude, W. G. Phosphoranyl Radicals - Their Structure, Formation, and Reactions. *Acc. Chem. Res.* **1982**, *15*, 117–125. (b) Bentrude, W. G.; Hansen, E. R.; Khan, W. A.; Min, T. B.; Rogers, P. E. Free-Radical Chemistry of Organophosphorus Compounds. III. α vs. β Scission in Reactions of Alkoxy and Thiyl Radicals with Trivalent Organophosphorus Derivatives. *J. Am. Chem. Soc.* **1973**, *95*, 2286–2293. (c) Bentrude, W. G.; Hansen, E. R.; Khan, W. A.; Rogers, P. E. α vs. β Scission in Reactions of Alkoxy and Thiyl Radicals with Diethyl alkylphosphonites. *J. Am. Chem. Soc.* **1972**, *94*, 2867–2868. (d) Davies, A. G.; Dennis, R. W.; Griller, D.; Roberts, B. P. A Kinetic Study of α - and β -scission of Phosphoranyl Radicals, Rn.P(OR)_{4-n}. *J. Organomet. Chem.* **1972**, *40*, 33–35. (e) Roberts, B. P.; Singh, K. Site-Selectivity in the β -Scission of Alkoxyphosphoranyl Radicals. A Reinterpretation. *J. Chem. Soc., Perkin Trans. 2* **1980**, 1549–1556.
- (13) (a) Zhang, L.; Koreeda, M. Radical Deoxygenation of Hydroxyl Groups via Phosphites. *J. Am. Chem. Soc.* **2004**, *126*, 13190–13191. (b) Ding, B.; Bentrude, W. G. Trimethyl Phosphite as a Trap for Alkoxy Radicals Formed from the Ring Opening of Oxiranylcarbonyl Radicals. Conversion to Alkenes. Mechanistic Applications to the Study of C–C versus C–O Ring Cleavage. *J. Am. Chem. Soc.* **2003**, *125*, 3248–3259. (c) Jordan, P. A.; Miller, S. J. An Approach to the Site-Selective Deoxygenation of Hydroxy Groups Based on Catalytic Phosphoramidite Transfer. *Angew. Chem., Int. Ed.* **2012**, *51*, 2907–2911.
- (14) (a) Rossi-Ashton, J. A.; Clarke, A. K.; Unsworth, W. P.; Taylor, R. J. K. Phosphoranyl Radical Fragmentation Reactions Driven by Photoredox Catalysis. *ACS Catal.* **2020**, *10*, 7250–7261. (b) Hu, X.-Q.; Hou, Y.-X.; Liu, Z.-K.; Gao, Y. Recent Advances in Phosphoranyl Radical-Mediated Deoxygenative Functionalisation. *Org. Chem. Front.* **2020**, *7*, 2319–2324. (c) Luo, K.; Yang, W. C.; Wu, L. Photoredox Catalysis in Organophosphorus Chemistry. *Asian J. Org. Chem.* **2017**, *6*, 350–367. (d) Pan, D.; Nie, G.; Jiang, S.; Li, T.; Jin, Z. Radical Reactions Promoted by Trivalent Tertiary Phosphines. *Org. Chem. Front.* **2020**, *7*, 2349–2371. (e) Shao, X.; Zheng, Y.; Ramadoss, V.; Tian, L.; Wang, Y. Recent Advances in P(III)-Assisted Deoxygenative Reactions Under Photochemical or Electrochemical Conditions. *Org. Biomol. Chem.* **2020**, *18*, 5994–6005.
- (15) (a) Han, J. B.; Guo, A.; Tang, X. Y. Alkylation of Allyl/Alkenyl Sulfones by Deoxygenation of Alkoxy Radicals. *Chem. - Eur. J.* **2019**, *25*, 2989–2994. (b) Gao, X. F.; Du, J. J.; Liu, Z.; Guo, J. Visible-Light-Induced Specific Desulfurization of Cysteiny Peptide and Glycopeptide in Aqueous Solution. *Org. Lett.* **2016**, *18*, 1166–1169. (c) Zheng, L.; Xia, P. J.; Zhao, Q. L.; Qian, Y. E.; Jiang, W. N.; Xiang, H. Y.; Yang, H. Photocatalytic Hydroacylation of Alkenes by Directly Using Acyl Oximes. *J. Org. Chem.* **2020**, *85*, 11989–11996. (d) Ye, Z.-P.; Hu, Y.-Z.; Xia, P.-J.; Xiang, H.-Y.; Chen, K.; Yang, H. Photocatalytic Intermolecular Anti-Markovnikov Hydroamination of Unactivated Alkenes with N-Hydroxyphthalimide. *Org. Chem. Front.* **2021**, *8*, 273–277.
- (16) (a) Cooper, J. W.; Roberts, B. P. Configurational Effects in the α -Scission of Phosphoranyl Radicals. *J. Chem. Soc., Perkin Trans. 2* **1976**, 808–813. (b) Trippett, S. Apicophilicity and Ring-Strain in Five-Coordinate Phosphoranes. *Phosphorus Sulfur Relat. Elem.* **1976**, *1*, 89–98.
- (17) (a) Koksai, Z.; Kalin, R.; Camadan, Y.; Usanmaz, H.; Almaz, Z.; Gulcin, I.; Gokcen, T.; Goren, A. C.; Ozdemir, H. Secondary Sulfonamides as Effective Lactoperoxidase Inhibitors. *Molecules* **2017**, *22*, 793–801. (b) Supuran, C. T. Special Issue: Sulfonamides. *Molecules* **2017**, *22*, 1642–1646. (c) Zhao, C.; Rakesh, K. P.; Ravidar, L.; Fang, W.-Y.; Qin, H.-L. Pharmaceutical and Medicinal Significance of Sulfur (SVI)-Containing Motifs for Drug Discovery: A Critical Review. *Eur. J. Med. Chem.* **2019**, *162*, 679–734. (d) Apaydin, S.; Török, M. Sulfonamide Derivatives as Multi-Target Agents for Complex Diseases. *Bioorg. Med. Chem. Lett.* **2019**, *29*, 2042–2050.
- (18) (a) Fei, J.; Wang, Z.; Cai, Z.; Sun, H.; Cheng, X. Synthesis of α -Tertiary Amine Derivatives by Intermolecular Hydroamination of Unfunctionalized Alkenes with Sulfamates under Trifluoromethanesulfonic Acid Catalysis. *Adv. Synth. Catal.* **2015**, *357*, 4063–4068. (b) Karshtedt, D.; Bell, A. T.; Tilley, T. D. Platinum-Based Catalysts for the Hydroamination of Olefins with Sulfonamides and Weakly Basic Anilines. *J. Am. Chem. Soc.* **2005**, *127*, 12640–12646. (c) Zhang, J.; Yang, C.-G.; He, C. Gold(I)-Catalyzed Intra- and Intermolecular Hydroamination of Unactivated Olefins. *J. Am. Chem. Soc.* **2006**, *128*, 1798–1799.
- (19) Nguyen, T. M.; Manohar, N.; Nicewicz, D. A. Anti-Markovnikov Hydroamination of Alkenes Catalyzed by a Two-Component Organic Photoredox System: Direct Access to Phenethylamine Derivatives. *Angew. Chem., Int. Ed.* **2014**, *53*, 6198–6201.
- (20) Zhu, Q.; Graff, D. E.; Knowles, R. R. Intermolecular Anti-Markovnikov Hydroamination of Unactivated Alkenes with Sulfonamides Enabled by Proton-Coupled Electron Transfer. *J. Am. Chem. Soc.* **2018**, *140*, 741–747.
- (21) (a) Qin, Q.; Ren, D.; Yu, S. Visible-Light-Promoted Chloramination of Olefins with N-Chlorosulfonamide as Both Nitrogen and Chlorine Sources. *Org. Biomol. Chem.* **2015**, *13*, 10295–10298. (b) Chen, J.; Guo, H. M.; Zhao, Q. Q.; Chen, J. R.; Xiao, W. J. Visible Light-Driven Photocatalytic Generation of Sulfonamidyl Radicals for Alkene Hydroamination of Unsaturated Sulfonamides. *Chem. Commun.* **2018**, *54*, 6780–6783. (c) Hu, X.-Q.; Chen, J.-R.; Wei, Q.; Liu, F.-L.; Deng, Q.-H.; Beauchemin, A. M.; Xiao, W.-J. Photocatalytic Generation of N-Centered Hydrazonyl Radicals: A Strategy for Hydroamination of β,γ -Unsaturated Hydrazones. *Angew. Chem., Int. Ed.* **2014**, *53*, 12163–12167. (d) Miyazawa, K.; Koike, T.; Akita, M. Regiospecific Intermolecular Aminohydroxylation of Olefins by Photoredox Catalysis. *Chem. - Eur. J.* **2015**, *21*, 11677–11680.
- (22) (a) Lardy, S. W.; Schmidt, V. A. Intermolecular Radical Mediated Anti-Markovnikov Alkene Hydroamination Using N-

Hydroxyphthalimide. *J. Am. Chem. Soc.* **2018**, *140*, 12318–12322. (b) Miller, D. C.; Ganley, J. M.; Musacchio, A. J.; Sherwood, T. C.; Ewing, W. R.; Knowles, R. R. Anti-Markovnikov Hydroamination of Unactivated Alkenes with Primary Alkyl Amines. *J. Am. Chem. Soc.* **2019**, *141*, 16590–16594. (c) Nguyen, S. T.; Zhu, Q.; Knowles, R. R. PCET-Enabled Olefin Hydroamidation Reactions with N-Alkyl Amides. *ACS Catal.* **2019**, *9*, 4502–4507.

(23) (a) Yasui, S.; Shioji, K.; Tsujimoto, M.; Ohno, A. Reactivity of a Trivalent Phosphorus Radical Cation as an Electrophile Toward Pyridine Derivatives. *Heteroat. Chem.* **2000**, *11*, 152–157. (b) Yasui, S.; Yamazaki, S. Intramolecular Stabilization of the Phosphine Radical Cation by the Second Phosphorus Atom during the Photooxidation of Diphosphines: 31P NMR Spectroscopic Analysis. *Chem. Lett.* **2015**, *44*, 422–424.

(24) (a) Zagumennov, V. A.; Sizova, N. A.; Nikitin, E. V. Electrochemical Oxidation of Tertiary Phosphines in the Presence of Camphene. *Russ. J. Gen. Chem.* **2009**, *79*, 1473–1482. (b) Ohmori, H.; Takanami, T.; Masui, M. Reaction of Triphenylphosphine Radical Cation with Cycloalkenes: Electrochemical One-Step Preparation of 1-Cycloalkenyltriphenylphosphonium Salts. *Tetrahedron Lett.* **1985**, *26*, 2199–2200.

(25) (a) Yasui, S.; Shioji, K.; Ohno, A. Reactivity of a Phosphoranyl Radicals Generated by Photoreaction of Phenyl Diphenylphosphinite with 10-Methylacridium Iodide - Alpha-Scission vs Single-Electron Transfer. *Heteroat. Chem.* **1994**, *5*, 85–90. (b) Yasui, S.; Shioji, K.; Tsujimoto, M.; Ohno, A. Kinetic Study on the Reaction of Tributylphosphine with Methylviologen. Reactivity of the Phosphine Radical Cation Intermediate Towards nucleophiles. *J. Chem. Soc., Perkin Trans. 2* **1999**, 855–862.

(26) Reactivity trends observed during optimization efforts were also observed in reactions conducted under optimized conditions. See section 2.2 of the [Supporting Information](#) for full details.

(27) Haav, K.; Saame, J.; Kütt, A.; Leito, I. Basicity of Phosphanes and Diphosphanes in Acetonitrile. *Eur. J. Org. Chem.* **2012**, *2012*, 2167–2172.

(28) A control reaction conducted by using N-methylbenzene-sulfonamide as a nucleophile showed no reactivity, confirming that second functionalization events are not possible under the reaction conditions.

(29) (a) Mayer, J. M. Proton-Coupled Electron Transfer: A Reaction Chemist's View. *Annu. Rev. Phys. Chem.* **2004**, *55*, 363–390. (b) Darcy, J. W.; Koronkiewicz, B.; Parada, G. A.; Mayer, J. M. A Continuum of Proton-Coupled Electron Transfer Reactivity. *Acc. Chem. Res.* **2018**, *51*, 2391–2399. (c) Weinberg, D. R.; Gagliardi, C. J.; Hull, J. F.; Murphy, C. F.; Kent, C. A.; Westlake, B. C.; Paul, A.; Ess, D. H.; McCafferty, D. G.; Meyer, T. J. Proton-Coupled Electron Transfer. *Chem. Rev.* **2012**, *112*, 4016–4093.

(30) Arias-Rotondo, D. M.; McCusker, J. K. The photophysics of photoredox catalysis: a roadmap for catalyst design. *Chem. Soc. Rev.* **2016**, *45*, 5803–5820.

(31) The absorption spectrum of the sulfonamide was not found to interfere with either the emitted wavelength of light for photoexcitation or the monitored wavelength of photocatalyst fluorescence.

(32) Warren, J. J.; Tronic, T. A.; Mayer, J. M. Thermochemistry of Proton-Coupled Electron Transfer Reagents and its Implications. *Chem. Rev.* **2010**, *110*, 6961–7001.

(33) Kargin, Y. M.; Budnikova, Y. G. Electrochemistry of Organophosphorus Compounds. *Russ. J. Gen. Chem.* **2001**, *71*, 1393–1421.

(34) Aminophosphine L-1 is only observed in the absence of olefin or in reactions with sterically bulky phosphines.

(35) (a) Hilton, M. C.; Zhang, X.; Boyle, B. T.; Alegre-Requena, J. V.; Paton, R. S.; McNally, A. Heterobiaryl Synthesis by Contractive C–C Coupling via P(V) Intermediates. *Science* **2018**, *362*, 799–804. (b) Hilton, M. C.; Dolewski, R. D.; McNally, A. Selective Functionalization of Pyridines via Heterocyclic Phosphonium Salts. *J. Am. Chem. Soc.* **2016**, *138*, 13806–13809. (c) Patel, C.; Mohnike, M.; Hilton, M. C.; McNally, A. A Strategy to Aminate Pyridines,

Diazines, and Pharmaceuticals via Heterocyclic Phosphonium Salts. *Org. Lett.* **2018**, *20*, 2607–2610.

(36) (a) Reichl, K. D.; Radosevich, A. T. A Phosphine-Mediated Stereocontrolled Synthesis of Z-Enediynes by a Vicinal Dialkynylation of Ethynylphosphonium Salts. *Chem. Commun.* **2014**, *50*, 9302–9305. (b) Uchida, Y.; Kozawa, H. Formation of 2,2'-Bipyridyl by Ligand Coupling on the Phosphorus Atom. *Tetrahedron Lett.* **1989**, *30*, 6365–6368.

(37) Parsaee, F.; Senarathna, M. C.; Kannangara, P. B.; Alexander, S. N.; Arche, P. D. E.; Welin, E. R. Radical Philicity and its Role in Selective Organic Transformations. *Nat. Rev. Chem.* **2021**, *5*, 486–499.

(38) Syed Shoaib Ahmad, S.; Gildardo, R.; Muhammad, A. Recent Advances in Medicinal Chemistry of Sulfonamides. Rational Design as Anti-Tumoral, Anti-Bacterial and Anti-Inflammatory Agents. *Mini-Rev. Med. Chem.* **2013**, *13*, 70–86.

(39) Wishart, D. S.; Feunang, Y. D.; Guo, A. C.; Lo, E. J.; Marcu, A.; Grant, J. R.; Sajed, T.; Johnson, D.; Li, C.; Sayeeda, Z.; Assempour, N.; Iynkkaran, I.; Liu, Y.; Maciejewski, A.; Gale, N.; Wilson, A.; Chin, L.; Cummings, R.; Le, D.; Pon, A.; Knox, C.; Wilson, M. DrugBank 5.0: A Major Update to the DrugBank Database for 2018. *Nucleic Acids Res.* **2018**, *46*, 1074–1082.